

Submitted to  
**Topology Proceedings**

## EFIMOV SPACES, CH, AND SIMPLE EXTENSIONS

ALAN DOW AND ROBERTO PICHARDO-MENDOZA

**ABSTRACT.** We give a construction under CH of an inverse system of simple extensions so that its limit is an Efimov space. This example shows that CH alone implies that a conjecture of Mercourakis about measures [8] is false.

### 1. INTRODUCTION

An Efimov space is an infinite compact Hausdorff space with no non-trivial converging sequences and no copies of  $\beta\omega$ , the Stone-Ćech compactification of the integers.

One of the well-known examples of this kind of space was constructed by Fedorčuk [5] with the aid of  $\diamond$ . His space is the limit of a special class of inverse system: continuous and based on simple extensions. The construction we present here is an improvement in the sense that we are just assuming CH.

Džamonja and Plebanek [3] show that any Efimov space constructed from the the Cantor space in an inverse limit of length  $\omega_1$  using simple extensions refutes a conjecture of Mercourakis concerning measures on compact spaces. Thus Fedorčuk's example does the job under  $\diamond$ . It is asked in [3] if CH suffices to refute Mercourakis' conjecture, and our construction answers this affirmatively.

---

1991 *Mathematics Subject Classification.* 54A25 (5AD30 04A15).

*Key words and phrases.* Efimov space, CH..

The first author acknowledges support provided by NSF grant DMS-20060114.

The second author was supported by the CONACYT grant no. 159538/215933.

Efimov spaces have been constructed from CH before (see, for example, [3] and [6]) but ours is the first example which is an inverse limit of simple extensions.

## 2. THE CONSTRUCTION

Recall that  $\langle f_{\alpha\beta}, X_\alpha : \alpha < \beta < \varepsilon \rangle$  is an inverse system if  $f_{\alpha\beta}$  is a map from  $X_\beta$  to  $X_\alpha$  whenever  $\alpha < \beta < \varepsilon$  and the equation  $f_{\alpha\gamma} = f_{\alpha\beta} \circ f_{\beta\gamma}$  holds for all  $\alpha < \beta < \gamma < \varepsilon$ . The inverse limit of the system is the appropriate subspace of the topological product of the family  $\{X_\alpha : \alpha < \varepsilon\}$  as described in [4].

**Definition 2.1.** An inverse system  $\langle f_{\alpha\beta}, X_\alpha : \alpha < \beta < \varepsilon \rangle$  is

- (1) *continuous* if  $X_\alpha$  is the inverse limit of the system  $\langle f_{\beta\gamma}, X_\beta : \beta < \gamma < \alpha \rangle$ , for any limit ordinal  $\alpha < \varepsilon$ .
- (2) *based on simple extensions* if  $X_{\alpha+1}$  is always a simple extension of  $X_\alpha$ , i.e. there exists a single point  $x_\alpha \in X_\alpha$  so that  $f_{\alpha, \alpha+1}^{-1}[x_\alpha]$  contains exactly two points and  $f_{\alpha, \alpha+1}^{-1}[x]$  is a singleton for all  $x \in X_\alpha \setminus \{x_\alpha\}$

These kinds of inverse systems are known as *minimal* in the Boolean algebra setting.

The following result, in Boolean algebraic form, is due to Koppelberg [7]; a topological proof can be found in [1].

**Proposition 2.2.** *If  $X$  is the limit of the inverse system of simple extensions  $\langle f_{\alpha\beta}, X_\alpha : \alpha < \beta < \varepsilon \rangle$ , then  $X$  does not map onto  $[0, 1]^{\omega_1}$ , unless  $X_0$  does.*

Observe that in this case,  $X$  does not map onto  $[0, 1]^c$  either and therefore  $\beta\omega$  cannot be embedded in  $X$ .

**Theorem 2.3.** *Under CH, there exists an Efimov space that can be obtained as the limit of an inverse system of simple extensions of length  $\omega_1$ .*

*Proof.* The plan is to use induction over  $\beta$  to get the desired inverse system  $\langle f_{\alpha\beta}, X_\alpha : \alpha < \beta < \omega_1 \rangle$ .

From now on, sequence means infinite sequence. Recall that a countable set  $E$  converges to a point  $a$  if  $E \setminus U$  is finite whenever  $U$  is a neighborhood of  $a$ .

Assume that we have constructed  $X_\alpha$  and  $X_\beta$  for some  $\alpha < \beta$  and let  $x \in X_\alpha$  be arbitrary. To simplify our notation we will write  $[x]_\beta$  to denote  $f_{\alpha\beta}^{-1}[\{x\}]$ .

Fix a partition  $\langle P_\alpha : \alpha < \omega_1 \rangle$  of  $\omega_1$  so that  $P_\alpha$  is an uncountable subset of  $\omega_1 \setminus \alpha$  for each  $\alpha < \omega_1$ .

We need some terminology. Let  $\gamma < \omega_1$  and  $\beta \in P_\gamma$  be arbitrary. If  $X_\gamma$  and  $X_\beta$  have been constructed, then use CH to fix an enumeration  $\langle D_\xi : \xi \in P_\gamma \rangle$  of all converging sequences  $D$  in  $X_\gamma$  so that  $f_{\xi\gamma}[D]$  is finite for each  $\xi < \gamma$ . Observe that  $D_\beta$  is a convergent sequence in  $X_\gamma$ . A *selector* from  $D_\beta$  is a set  $E \subseteq X_\beta$  so that  $E \cap [d]_\beta$  is a singleton for each  $d \in D_\beta$ . Use CH again to get  $\langle E_\xi : \xi \in P_\beta \rangle$ , an enumeration of all converging selectors from  $D_\beta$ .

Let  $\beta < \alpha < \varepsilon$  and  $x \in X_\alpha$  be arbitrary. Fix  $\gamma < \omega_1$  so that  $\beta \in P_\gamma$ . We define

$$D_\beta(x) := \{d \in D_\beta : x \notin [d]_\alpha\}.$$

In other words,  $d \in D_\beta(x)$  if and only if  $d \in D_\beta$  and  $f_{\gamma\alpha}(x) \neq d$ .

Assume that for some  $\varepsilon < \omega_1$  we have defined a continuous inverse system of simple extensions  $\langle f_{\gamma\beta}, X_\gamma : \gamma < \beta < \varepsilon \rangle$  so that the following holds for each  $\beta < \varepsilon$ .

- (1)  $X_0 = 2^\omega$ , the Cantor set.
- (2) If  $\beta + 1 < \varepsilon$  and  $i < 2$ , then there exist  $A_\beta^i$ , a closed subset of  $X_\beta$ , and  $H_\beta^i$ , an infinite subset of  $E_\beta$ , so that
  - (a)  $A_\beta^0 \cap A_\beta^1 = \{x_\beta\}$ ,
  - (b)  $X_\beta = A_\beta^0 \cup A_\beta^1$ ,
  - (c)  $X_{\beta+1} = A_\beta^0 \oplus A_\beta^1$  (the topological sum), and
  - (d)  $f_{\beta,\beta+1}$  is the projection map.
  - (e)  $[e]_\beta \subseteq A_\beta^i$  for all  $e \in H_\beta^i$ .
- (3) If  $x \in U \in \mathcal{T}_\beta$ , where  $\mathcal{T}_\beta$  is the topology of  $X_\beta$ , then for each finite set  $F \subseteq \beta$  there exists  $W$ , a clopen subset of  $X_\beta$ , such that  $x \in W \subseteq U$  and  $W$  takes care of  $(x, F)$ , i.e.

$$[d]_\gamma \cap f_{\gamma\beta}[W] \cap f_{\gamma\beta}[X_\beta \setminus W] = \emptyset,$$

for all  $\gamma \in F$  and  $d \in D_\gamma(x)$ .

Condition (3) is equivalent to saying that  $W \cap [d]_\beta$  is a preimage of a clopen subset of  $[d]_\gamma$ .

Observe that (1) guarantees, according to Proposition 2.2, that the limit will not contain a copy of  $\beta\omega$ . Hence our main concern is to get rid of all converging sequences.

If  $\varepsilon$  is a limit ordinal, then  $X_\varepsilon$  is the limit of this inverse system. To verify (3) let  $x \in U_0 \in \mathcal{T}_\varepsilon$  be arbitrary. Fix a finite set  $F \subseteq \varepsilon$ . Since  $\{f_{\xi\varepsilon}^{-1}[V] : \xi < \varepsilon, V \in \mathcal{T}_\xi\}$  is a base for  $\mathcal{T}_\varepsilon$ , there exists  $\alpha < \varepsilon$  and  $U \in \mathcal{T}_\alpha$  so that  $F \subseteq \alpha$  and  $x \in f_{\alpha\varepsilon}^{-1}[U] \subseteq U_0$ . Apply the inductive hypothesis to  $\alpha$ ,  $f_{\alpha\varepsilon}(x)$ ,  $U$ , and  $F$  to get a clopen set  $W$  in  $X_\alpha$  which takes care of  $(f_{\alpha\varepsilon}(x), F)$  and satisfies  $f_{\alpha\varepsilon}(x) \in W \subseteq U$ . For each  $\gamma \in F$  we have  $D_\gamma(x) = D_\gamma(f_{\alpha\varepsilon}(x))$ , so  $f_{\alpha\varepsilon}^{-1}[W]$  is a clopen subset of  $X_\varepsilon$  which takes care of  $(x, F)$ .

Now assume that  $\varepsilon = \alpha + 1$ . Let  $\beta < \omega_1$  be such that  $\alpha \in P_\beta$ . Hence  $\beta \leq \alpha$  and therefore  $E_\alpha$  and  $D_\beta$  have been defined. Since  $X_0$  is compact metrizable and  $\alpha < \omega_1$ ,  $X_\alpha$  is compact metrizable too. In particular the family  $\{[e]_\alpha : e \in E_\alpha\}$  must have an accumulation point, i.e. there exists a point  $x_\alpha$  in  $X_\alpha$  so that the set  $\{e \in E_\alpha : [e]_\alpha \cap V \neq \emptyset\}$  is infinite for each neighborhood  $V$  of  $x_\alpha$ .

The next step is to find an infinite set  $H_\alpha \subseteq E_\alpha$  so that  $\langle [e]_\alpha : e \in H_\alpha \rangle$  converges to  $x_\alpha$ , i.e. for each neighborhood  $U$  of  $x_\alpha$  all but finitely many  $e \in H_\alpha$  satisfy  $[e]_\alpha \subseteq U$ . Let's start by fixing a local decreasing base  $\{B_n : n \in \omega\}$  for  $x_\alpha$ .

We face two cases. When  $\beta < \alpha$  apply (3) to construct a sequence  $\langle W_n : n \in \omega \rangle$  of clopen subsets of  $X_\alpha$  which satisfies

- (i)  $x_\alpha \in W_0 \subseteq B_0$ ,
- (ii)  $x_\alpha \in W_{n+1} \subseteq W_n \cap B_{n+1}$ , and
- (iii)  $W_n$  takes care of  $(x_\alpha, \{\beta\})$ ,

for each  $n \in \omega$ .

Define  $E_\alpha^n := \{e \in E_\alpha : [e]_\alpha \cap W_n \neq \emptyset\}$ . We claim that for all but possibly one  $e \in E_\alpha^n$  we get  $[e]_\alpha \subseteq W_n$ . To prove this assertion let  $e \in E_\alpha^n$  and  $d \in D_\beta(x_\alpha)$  be so that  $e \in [d]_\beta$ . Now note that if  $[e]_\alpha \setminus W_n \neq \emptyset$ , then  $e \in f_{\beta\alpha}[W_n] \cap f_{\beta\alpha}[X_\alpha \setminus W_n] \cap [d]_\beta$ , a clear contradiction to (iii). Find an infinite set  $H_\alpha \subseteq E_\alpha$  so that  $H_\alpha \setminus E_\alpha^n$  is finite for all  $n \in \omega$  and observe that this  $H_\alpha$  works.

For the case  $\alpha = \beta$  we have  $E_\alpha \subseteq X_\alpha$  and  $[e]_\alpha = \{e\}$ , for each  $e \in E_\alpha$ . Clearly any subsequence of  $E_\alpha$  which converges to  $x_\alpha$  will work as  $H_\alpha$ .

Let  $\alpha = \{\beta_k : k < \omega\}$  be an enumeration of  $\alpha$ . Use (3) to construct  $\{e_n : n \in \omega\} \subseteq H_\alpha$ ,  $g : \omega \rightarrow \omega$ , and  $\{U_n : n \in \omega\}$  so that for each  $n \in \omega$

- (I)  $U_0 = X_\alpha$ ,
- (II)  $U_n$  is clopen in  $X_\alpha$  and takes care of  $(x_\alpha, \{\beta_k : k \leq n\})$ ,
- (III)  $g$  is an increasing function,
- (IV)  $x_\alpha \in U_{n+1} \subseteq B_{g(n)} \subseteq U_n \setminus [e_n]_\alpha$ , and
- (V)  $[e_n]_\alpha \subseteq U_n$ .

We are going to partition  $X_\alpha \setminus \{x_\alpha\}$ . Let  $n$  be an arbitrary integer. Observe that the set  $V_n := U_n \setminus U_{n+1}$  is clopen and takes care of  $(x_\alpha, \{\beta_k : k \leq n\})$ . Now, given  $i < 2$ , define

$$b_n^i := \{x_\alpha\} \cup \bigcup_{k=n}^{\infty} V_{2k+i}.$$

The following holds for each  $i < 2$ .

- (a)  $b_n^i$  is closed for all  $n < \omega$ .
- (b) If  $U$  a neighborhood of  $x_\alpha$ , then  $b_m^i \subseteq U$ , for some  $m < \omega$ .

We claim that  $b_n^i$  takes care of  $(x_\alpha, \{\beta_k : k \leq n\})$ . Let  $k \leq n$  and  $d \in D_{\beta_k}(x_\alpha)$  be arbitrary. If  $y \in f_{\beta_k \alpha}[b_n^i] \cap f_{\beta_k \alpha}[X_\alpha \setminus b_n^i]$ , then we have two possibilities:  $y \in f_{\beta_k \alpha}[V_{2\ell+i}]$ , for some  $\ell \geq n$ , or  $y = f_{\beta_k \alpha}(x_\alpha)$ . In the first case we get  $y \in f_{\beta_k \alpha}[V_{2\ell+i}] \cap f_{\beta_k \alpha}[X_\alpha \setminus V_{2\ell+i}]$  and  $k \leq n \leq \ell \leq 2\ell + i$ , so  $y \notin [d]_{\beta_k}$ .

Now assume that  $y = f_{\beta_k \alpha}(x_\alpha)$ . Since  $d \in D_{\beta_k}(x_\alpha)$ , we get  $x_\alpha \notin [d]_\alpha$  and therefore  $y \notin [d]_{\beta_k}$ .

For each  $i < 2$ , set  $A_\alpha^i := b_0^i$  and  $H_\alpha^i := \{e_{2n+i} : n \in \omega\}$ . Let  $X_{\alpha+1} := (A_\alpha^0 \times \{0\}) \cup (A_\alpha^1 \times \{1\})$  and declare open all the sets of the form  $(U^0 \times \{0\}) \cup (U^1 \times \{1\})$ , where  $U^i$  is open in the subspace topology of  $A_\alpha^i \subseteq X_\alpha$ . The map  $f_{\alpha, \alpha+1} : X_{\alpha+1} \rightarrow X_\alpha$  is defined by  $f_{\alpha, \alpha+1}(x, i) = x$ .

To complete the induction we check that property (3) holds. Assume that  $(x, i) \in U \in \mathcal{T}_{\alpha+1}$  are arbitrary and fix a finite set  $F \subseteq \alpha + 1$ . If  $x \neq x_\alpha$ , find  $U_0 \in \mathcal{T}_\alpha$  so that  $(x, i) \in U_0 \times \{i\} \subseteq U$  and  $x_\alpha \notin U_0$ . Let  $W_0$  be a clopen subset of  $X_\alpha$  which takes care of  $(x, F \setminus \{\alpha\})$  and satisfies  $x \in W_0 \subseteq U_0$ . Set  $W := W_0 \times \{i\}$  and note that for all  $\beta \in F$  we have  $f_{\beta, \alpha+1}[W] = f_{\beta \alpha}[W_0]$  and  $f_{\beta, \alpha+1}[X_{\alpha+1} \setminus W] = f_{\beta \alpha}[X_\alpha \setminus W_0]$  (recall that  $f_{\alpha \alpha}$  is the identity map). Since  $D_\beta((x, i)) = D_\beta(x)$  we conclude that  $W$  takes care of  $((x, i), F)$ .

Assume now that  $x = x_\alpha$ . Find  $n \in \omega$ , so that  $b_n^i \times \{i\} \subseteq U$  and  $F \setminus \{\alpha\} \subseteq \{\beta_k : k \leq n\}$ . Define  $W := b_n^i \times \{i\}$ . For each  $\beta \in F$  and  $d \in D_\beta((x_\alpha, i))$  we have that  $f_{\beta\alpha}(x_\alpha) \notin [d]_\beta$  and

$$f_{\beta,\alpha+1}[W] \cap f_{\beta,\alpha+1}[X_{\alpha+1} \setminus W] = \{f_{\beta\alpha}(x_\alpha)\} \cup (f_{\beta\alpha}[b_n^i] \cap f_{\beta\alpha}[X_\alpha \setminus b_n^i]).$$

Therefore  $W$  takes care of  $((x_\alpha, i), F)$ .

Let  $X$  be the limit of our inverse system and let  $\pi_\alpha : X \rightarrow X_\alpha$  be the bonding map for each  $\alpha < \omega_1$ . In order to check that  $X$  is an Efimov space, assume that  $S$  is a converging sequence in  $X$ . Let  $\gamma < \omega_1$  be the least ordinal so that  $\pi_\gamma[S]$  is infinite. Since  $\pi_\gamma[S]$  is a convergent sequence in  $X_\gamma$ , there exists  $\beta \in P_\gamma$  so that  $\pi_\gamma[S] = D_\beta$ . Since  $f_{\gamma\beta} \circ \pi_\beta = \pi_\gamma$ , we can find an infinite set  $S_0 \subseteq S$  so that  $\pi_\beta$  is one-to-one on  $S_0$  and  $\pi_\beta[S_0]$  is a selector from  $D_\beta$ , i.e. there exists  $\alpha \in P_\beta$  so that  $\pi_\beta[S_0] = E_\alpha$ . Property (2) provides two infinite subsets of  $S_0$ , namely  $S_0^0$  and  $S_0^1$ , so that  $\pi_\alpha[S_0^i] \subseteq A_\alpha^i$  for each  $i < 2$ . Therefore  $\pi_{\alpha+1}[S_0^0]$  and  $\pi_{\alpha+1}[S_0^1]$  cannot converge to the same point. This contradiction ends the proof.  $\square$

#### REFERENCES

- [1] A. Dow, *Efimov spaces and the splitting number*. Spring Topology and Dynamical Systems Conference. Topology Proc. **29** (2005), no. 1, 105–113.
- [2] A. Dow, D.H. Fremlin, *Compact sets without converging sequences in the random real model*, Acta Math. Univ. Comenian. (N.S.) **76** (2007), no. 2, 161–171. 54A25
- [3] M. Džamonja, G. Plebanek, *On Efimov spaces and Radon measures*, Topology Appl. **154** (2007), no. 10, 2063–2072.
- [4] R. Engelking, *General topology*, second ed., Sigma Series in Pure Mathematics, vol. 6, Heldermann Verlag, Berlin, 1989, Translated from the Polish by the author. MR **91c**:54001
- [5] V. V. Fedorčuk, *Completely closed mappings, and the consistency of certain general topology theorems with the axioms of set theory*, Matematicheskii Sbornik. Novaya Seriya **99** (**141**) (1976), 1–26 (Russian); English transl. Mathematics of the USSR. Sbornik **28** (1976), 3–33, 135. MR **53** #14379, Zbl. 0371.54045
- [6] K.P. Hart, *Efimov's problem*, Open Problems in Topology II, (E. Pearl, ed.), Elsevier Science, 2007, pp. 171–177.
- [7] S. Koppelberg, *Minimally generated Boolean algebras*, Order **5** (1989), no. 4, 393–406. MR **90g**:06022
- [8] S. Mercourakis, *Some remarks on countably determined measures and uniform distribution of sequences*, Monats. Math. **121** (1996), 79–101.

DEPARTMENT OF MATHEMATICS, UNC-CHARLOTTE, 9201 UNIVERSITY CITY  
BLVD., CHARLOTTE, NC 28223-0001

*E-mail address:* `adow@uncc.edu`

DEPARTMENT OF MATHEMATICS, UNC-CHARLOTTE, 9201 UNIVERSITY CITY  
BLVD., CHARLOTTE, NC 28223-0001

*E-mail address:* `rpichard@uncc.edu`