

## Assignment 11

### Oral questions


1. Assume  $a, b, c \in \mathbb{R}$  satisfy  $a^2 + bc = 1$ , and let  $T : \mathbb{C} \rightarrow \mathbb{C}$  be given by

$$T(z) = \frac{a\bar{z} + b}{c\bar{z} - a}.$$

Show that  $T(T(z)) = z$  for all  $z$ . (All reflections of the Poincaré upper half plane model are represented by such a function.)

2. Schweikart's constant is the distance  $d$  for which the angle of parallelism is  $\Pi(d) = 45^\circ$ . Prove that for the length function of the Poincaré disk model, Schweikart's constant equals  $\log(1 + \sqrt{2})$ . You may use the following formula in your proof. If a point  $P$  is at a Euclidean distance  $r$  from the center  $O$  then its hyperbolic distance from  $O$  is

$$d(O, P) = \ln \left( \frac{1+r}{1-r} \right).$$



### Questions to be answered in writing

1. Let  $a, b, c, d$  be real numbers, such that  $ad - bc \neq 0$ . Using that

$$\frac{az + b}{cz + d} = \begin{cases} \frac{a}{c} + \frac{b-ad/c}{cz+d} & \text{if } c \neq 0, \text{ and} \\ \frac{az+b}{d} & \text{if } c = 0, \end{cases}$$

show that every fractional linear transformation of the above form arises as a combination of horizontal translations  $z \mapsto z + b$ , dilations  $z \mapsto az$  and “reflected inversions”  $z \mapsto 1/z$ . Conclude that fractional linear transformations preserve angles and the cross-ratio.

2. A hyperbolic circle centered at  $C$  of radius  $r$  is the set of all points  $A$  satisfying  $d(A, C) = r$ . Prove that a hyperbolic circle corresponds to a circle in the Poincaré disk model, although its center may not be equal to its Euclidean center. (Hint: prove the statement in the case when  $C = P$  first, where  $P$  is the center of the Poincaré disk. Use then the fact that reflection about the perpendicular bisector of  $PC$  takes a hyperbolic circle centered at  $P$  into a hyperbolic circle centered at  $C$ , and that this reflection corresponds to an inversion about a circle.)