

May 6, 2002

Your name _____

As usual, show all your work. If you used a calculator, explain in detail how you used it to solve the problem. Part A is worth 120 points and part B 130 points.

1. (40 points) Find the following antiderivatives.

(a) $\int 6x^3 - 5x - 1 dx$

(b) $\int 6x^{\frac{3}{2}} + x^{-\frac{1}{2}} dx$

(c) $\int \frac{3x^3 + 2x - 1}{x} dx$

(d) $\int \frac{2x + 1}{x^2 + x - 3} dx$

(e) $\int 5x^4(x^5 + 2)^7 dx$

2. (20 points) Compute the following definite integrals.

(a) $\int_0^2 2xe^{-x^2} dx$

(b) $\int_0^5 (2x - 1)\sqrt{x^2 - x + 5} dx$

3. (15 points) Find a function $G(x)$ whose derivative is $3x^2 - 7$ and for which $G(4) = 9$.
4. (15 points) Find the area of the region bounded by $y = x^{3/2}$, the x -axis, and the lines $x = 0$ and $x = 4$.
5. (30 points) An object is thrown upwards from the top of a 400 feet high building, after which its path is governed entirely by gravity. The acceleration due to gravity is given by $a(t) = -32$, where t is measured in seconds and $a(t)$ is measured in ft/sec^2 . Recall that $a(t) = v'(t) = s''(t)$, where $v(t)$ denotes the velocity of the object (negative when its moving towards the earth), and $s(t)$ is the position of the object at time t . The equation above simply says that the velocity is an antiderivative of acceleration, and that position is an antiderivative of velocity. Suppose the initial velocity is 80 feet per second; ie, $v(0) = 80$. Answer the following questions about the path of the object.

- (a) Compute the function $v(t)$.

(b) Compute the function $s(t)$.

(c) At what time does the object hit the ground?

(d) At what time does the object reach its maximum height?

(e) What is its maximum height?

Part B

- (5 points) The slope of the line that contains the points $(-1, y)$ and $(4, -12)$ is -3 ? What is y ?
- (5 points) What is the slope of the line perpendicular to the line $2y + x = 4$?
- (10 points) Over what intervals is the second derivative of $g(x) = x^4 - 6x^3 + 12x^2 + 2x + 2$ negative?
- (15 points) Construct a cubic polynomial $f(x)$ that has zeros at $x = -2$, $x = 1$, and $x = 3$ and satisfies $f(0) = -12$.
- (10 points) Let $g(x)$ be defined as follows: Let

$$g(x) = \begin{cases} e^x & \text{if } x \leq 1 \\ \ln(x) & \text{if } x > 1 \end{cases}$$

Find an equation for the line tangent to the graph of $g(x)$ at the point $(3, g(3))$.

- (20 points) Compute each of the following derivatives.
 - $\frac{d}{dx} \frac{\sqrt{x^2-3}}{2x}$
 - $\frac{d}{dx} e^{x+\ln(x)}$
 - $\frac{d}{dx} \ln(x^2 + e^{2x})$
- (10 points) Calculate the doubling time for a 7% investment compounded continuously.
- (30 points) Suppose $u(x)$ is a function whose derivative $u'(x) = (2x + 1)^2(x - 2)^2$. Recall that theorem B tells you the intervals over which $u(x)$ is concave upwards based on $u''(x)$.
 - Compute $u''(x)$.
 - Find the three zeros of $u''(x)$.
 - Use the Test Interval Technique to find the intervals over which $u(x)$ is concave up.
- (5 points) What is $\lim_{h \rightarrow 0} \frac{1 - \sqrt{1 + 2h}}{h}$?
- (20 points) Find the absolute maximum and absolute minimum of the function $h(x) = \sqrt{x^2 + 6x + 25}$ over the interval $[-5, 5]$.